

# Impact of the Particle Environment on LYRA Data


M. Dominique, A. BenMoussa, M.Kruglanski, L. Dolla,  
I. Dammasch, M. Kretzschmar  
PROBA2 workshop, May 04 2012, Brussels




# PROBA2: Project for On-Board Autonomy

## PROBA2 orbit:



- ☐ Heliosynchronous
- ☐ Polar
- ☐ Dawn-dusk
- ☐ 725 km altitude
- ☐ Duration of 100 min



- ◆ Crosses the SAA about 8 times a day
- ◆ Crosses the auroral oval 4 times an orbit



launched on November 2, 2009



|       | Ly            | Hz           | Al                 | Zr                |
|-------|---------------|--------------|--------------------|-------------------|
|       | 120-123 nm    | 190-222 nm   | 17-80 nm +<br><5nm | 6-20 nm +<br><2nm |
| Unit1 | MSM - diamond | PIN- diamond | MSM- diamond       | P-N Silicon       |
| Unit2 | MSM- diamond  | PIN- diamond | MSM- diamond       | MSM- diamond      |
| Unit3 | P-N Silicon   | PIN- diamond | P-N Silicon        | P-N Silicon       |


# South Atlantic Anomaly





# SAA: effect on LYRA

**In 2010**

**Cover closed:**




**Covers open:**





# SAA: effect on LYRA


**In 2012**



- ☐ Effect of SAA constant
- ☐ Overall responsivity decreased (ageing)

=> SAA now visible in MSM diamond detectors of the nominal unit

**SAA produces peaks of amplitude equivalent to a B2 flare in unit 2**



B2 flare




- ❑ Independent on the pointing direction and on the covers status
- ❑ Independent on the spectral range
- ❑ Absolute amplitude of perturbation constant over the mission ( $\sim 0.5$  counts/ms in Si,  $\sim 0.05$  counts/ms in MSM diamond)
- ❑ Dependent on the detector material/type

| SWAP | LYRA        |                 |    |
|------|-------------|-----------------|----|
|      | Diamond PIN | Diamond MSM     | Si |
| ✓    | X           | Low sensitivity | ✓  |







2010-08-02 WDWSZE:02



Color representing max deviation at a point. 0.001 -> 5

## NASA AP-8/AE-8 Trapped radiation particle flux (SPENVIS)

**Protons > 20MeV**      **Electrons > 0.4 MeV**








## Energy deposition due to energetic protons

The surrounding shielding causes:

- ❑ slowdown the protons
- ❑ generation of secondary electrons



Collected in the bulk of the detector material

Energy needed to create 1 electron-hole pair is

- ❑ 1.1eV for Silicon
- ❑ 5.5 eV for diamond

Collected in surface:



- ❑ PIN diamond is not sensitive
- ❑ MSM diamond (planar structure) is slightly more sensitive
- ❑ PIN silicon is very sensitive.



# Energy deposition due to energetic protons

The surrounding shielding causes:


- slowdown the protons
- generation of secondary electrons



Energy deposition in Si behind a spherical Al shielding  
SHIELDOSE-2 (SPENVIS)




# LYRA's filters (Hz) after proton tests (@14.5MeV)



**Acton filters**

**Remark: same observation for Ly-a filters (XN and N)**


**After more than 2 years in orbit  
→ acc. fluence 7.1E9 (>10MeV)**






# Si detector (AXUV) after proton tests (@14.5MeV)

**NUV-VIS spectral response decreases (factor 1.5)**




**Dark current increases (x100)**






# Diamond detectors after proton tests (@14.5MeV)

## Dark current MSM24r




## Dark current (PIN11)

DC increases (x7) but still negligible  
(> pA @ 0V)





→ spectral response to be measured (soon)

# Perturbations in the auroral zone




# Auroral Oval

- ☐ Perturbations appearing around 75° latitude
- ☐ 2-3 days after a CME, flare ...
- ☐ Associated to geomagnetic perturbations of  $K_p \geq 4$
- ☐ Only in Al and Zr channels
- ☐ Seems to be sensitive to the ageing of the channel
- ☐ Not seen with covers closed








**Channel 3 in units 2 lost 95% of its sensitivity**

**BUT**

**The perturbations in channel 2 amplified by a factor 20 do not appear 20 X bigger than in channel 3.**

**=> The perturbation amplitude might be affected by the channel degradation**




## Possible origins of the auroral effect


- ☐ Galactic Cosmic Rays
- ☐ Protons or ions ejected by the Sun (SEP)
- ☐ Highly energetic electrons
- ☐ Photons
- ☐ ???



- ❑ The region in which the GCR are sensed is slightly wider after a geomagnetic storm, but it exists all the time
- ❑ GRC should be detected all over the polar caps



Incompatible with the zero-detection under normal geomagnetic conditions



## Possible origins of the auroral effect



- ☐ Galactic Cosmic ~~R~~ays
- ☐ Protons or ions ejected by the Sun (SEP)
- ☐ Highly energetic electrons
- ☐ Photons
- ☐ ???



Simulation with  
Magnetocosmics (SPENVIS):  
protons from outside the  
magnetosphere should be able  
to reach the altitude of the  
spacecraft for energy  $> 30$  MeV

BUT


The occurrence of SEP is not  
always correlated with the  
auroral perturbations observed  
by LYRA





## Possible origins of the auroral effect

- ☐ Galactic Co~~x~~mic Rays
- ☐ Protons or ion~~s~~ ejected by the Sun (SEP)
- ☐ Highly energetic electrons
- ☐ Photons
- ☐ ???



## Highly energetic electrons

- ☐ stopped by shielding  
except in the line of sight OK
- ☐ not seen by SWAP because of its off-  
line axis configuration OK
- ☐ only seen in Al and Zr => only ?  
explained if stopped by the thick  
interferential filters ( $\sim 7\text{mm}$ ) and not by  
the metallic ones (Al =  $158\text{nm}$  & Zr =  
 $148$  or  $300\text{nm}$ )
- ☐ ageing effects unexplained Non OK





## Possible origins of the auroral effect

- ☐ Galactic Co~~x~~mic Rays
- ☐ Protons or ion~~s~~ ejected by the Sun (SEP)
- ☐ Highly ener~~g~~getic electrons
- ☐ Photons
- ☐ ???




## ☐ Auroral:

- ☐ O+ line at 53.9 nm
- ☐ emission in the F layer, mostly below the altitude of PROBA2

## ☐ Airglow:

- ☐ He+ 30.4-nm, He 58.4-nm, O+ 53.9-nm
- ☐ emission region up to 1.25 ER


## ☐ Others?



From Sandel, B. R., et al.,  
Space Sci. Rev., 109, 25, 2003.)




# Filter + detector responsivity





- ☐ Auroral: **Too low altitudes**
  - ☐ O+ line at 53.9 nm
  - ☐ emission in the F layer, mostly below the altitude of PROBA2
- ☐ Airglow: **In auroral zones only**
  - ☐ He+ 30.4-nm, He 58.4-nm, O+ 53.9-nm
  - ☐ emission region up to 1.25 ER
- ☐ Others?




From Sandel, B. R., et al.,  
Space Sci. Rev., 109, 25, 2003.)




# Al vs Zr in unit 2 (degraded)

09/03/2012

Unit 2 – ch 3




Unit 2 – ch 4





# Al vs Zr in unit 2 (degraded)


LYRA timeseries, lev1, unit2



**In unit 2 (degraded unit), Al and Zr are identical**

**=> SXR photons?**

LYRA timeseries, lev1, unit2







# Aurora in AI channel

09/03/2012


Unit 2 – ch 3




Unit 3 – ch 3








**The amplitude of the auroral perturbation is more important in unit 3 (Si detectors, low degradation) than in unit 2 (diamond detectors, high degradation)**







# Aurora in Zr channel



**Again, perturbation in unit 3  
> in unit 2**

**=> Do we see EUV photons  
in the less degraded unit?**






## Possible origins of the auroral effect

- ☐ Galactic Co~~x~~mic Rays
- ☐ Protons or ion~~s~~ ejected by the Sun (SEP)
- ☐ Highly ener~~g~~etic electrons
- ☐ Photons ?
- ☐ ???



|                                     | GCR | SEP | Electrons | EUV<br>Photons | Others<br>(Brems-<br>strahlung ?) |
|-------------------------------------|-----|-----|-----------|----------------|-----------------------------------|
| Covers open only                    | ?   | ?   | V         | V              | ?                                 |
| In auroral zone                     | X   | V   | V         | V              | ?                                 |
| After major solar event             | X   | V   | V         | V              | ?                                 |
| In Al and Zr only                   | X   | X   | ?         | V              | ?                                 |
| ageing effect                       | X   | X   | X         | V              | ?                                 |
| Al and Zr of same amplitude in 2012 | ?   | X   | V         | V              | ?                                 |

X = incompatible  
V = compatible



## Conclusions

- ❑ The underlying process is still not clear to us. Both SWAP and LYRA sense energetic trapped protons in SAA
- ❑ LYRA senses an auroral signature in its two shorter wavelength channels.
- ❑ Work still in progress ...



<http://proba2.sidc.be/>



**European Space Agency**



**Belgian Science Policy Office**



The research leading to these results has received funding from the European Commission's Seventh Framework Programme (FP7/2007-2013) under the grant agreement eHeroes (project n° 284461, [www.eheroes.eu](http://www.eheroes.eu)).